

Pair production of the T-odd leptons at the LHC

Chong-Xing Yue, Yong-Zhi Wang, Wei Lui, Wei Ma

Department of Physics, Liaoning Normal University, Dalian 116029, P. R. China *

April 30, 2009

Abstract

The T-odd leptons predicted by the littlest *Higgs* model with T-parity can be pair produced via the subprocesses $gg \rightarrow \ell_H^+ \ell_H^-$, $q\bar{q} \rightarrow \ell_H^+ \ell_H^-$, $\gamma\gamma \rightarrow \ell_H^+ \ell_H^-$ and $VV \rightarrow \ell_H^+ \ell_H^-$ ($V=W$ or Z) at the *CERN* Large Hadron Collider (*LHC*). We estimate the hadronic production cross sections for all of these processes and give a simply phenomenology analysis. We find that the cross sections for most of the above processes are very small. However, the value of the cross section for the *Drell – Yan* process $q\bar{q} \rightarrow \ell_H^+ \ell_H^-$ can reach $270fb$.

PACS numbers: 12.60.Cn, 14.60.Hi, 13.85.Lg

*E-mail:cxyue@lnnu.edu.cn

1. Introduction

The *CERN* Large Hadron Collider (*LHC*) will soon go into full operation and provide proton-proton collisions at a center-of-mass (*c.m.*) energy of 14TeV , a factor of 7 higher than the Tevatron. There are strong theoretical reasons to expect that the *LHC* will discover new physics beyond the standard model (*SM*) up to TeV . Many popular new physics models beyond the *SM* predict the existence of the new charged leptons which are denoted as $L^\pm$. In general, as long as they are not too heavy, these new particles can be pair produced via the gluon fusion process $gg \rightarrow L^+L^-$, the *Drell – Yan* process $q\bar{q} \rightarrow L^+L^-$, the photon-induced process $\gamma\gamma \rightarrow L^+L^-$, and the weak gauge boson fusion process $VV \rightarrow L^+L^-$ ($V=W$ or Z) at the *LHC*.

Studying production and decay of the new charged leptons at current or in future high energy collider experiments is of special interest. Any signal for such kind of new fermions in future high energy experiments will play an important role in testing the *SM* flavor structure and discovery of new physics beyond the *SM*. This fact has lead to many works involving the new charged leptons at e^+e^- colliders [1], ep colliders [2], and hadron colliders [3,4,5].

The littlest *Higgs* model with T-parity, which is called the *LHT* model [6], is one of the attractive little *Higgs* models. The *LHT* model predicts the existence of the T-odd $SU(2)$ doublet fermions. These new fermions can produce rich phenomenology at present or in future high energy collider experiments [7,8,9,10,11]. In Ref.[12], we have studied pair production of the T-odd leptons in an international linear e^+e^- collider (*ILC*). Our numerical results show that, in wide range of the parameter space, they can be copiously produced in pairs. The possible signatures of the T-odd leptons might be detected in future *ILC* experiments.

Since it is regarded as the cross section for pair production of the T-odd leptons is generally small at the *LHC*, so far there are few works involved their directly production at the *LHC*. However, considering the *LHC* will go into operation and it is possible to completely study the possible signals of the T-odd leptons at the *LHC*, it is need to carefully study pair production of the T-odd leptons at the *LHC*. So, in this note, we will

discuss various possible production channels of the T-odd lepton pairs at the *LHC* and compare their values of the production cross sections for different production channels. In the next section, we will give our numerical results. Our conclusion and a simply phenomenology analysis are given in the last section.

2. The numerical results

The *LHT* model [6] is based on an $SU(5)/SO(5)$ global symmetry breaking pattern. A subgroup $[SU(2) \times U(1)]_1 \times [SU(2) \times U(1)]_2$ of the $SU(5)$ global symmetry is gauged, and at the scale f it is broken into the *SM* electroweak symmetry $SU(2)_L \times U(1)_Y$. T-parity is an automorphism that exchanges the $[SU(2) \times U(1)]_1$ and $[SU(2) \times U(1)]_2$ gauge symmetries. To simultaneously implement T-parity, one needs to double the *SM* fermion doublet spectrum [6,7]. The T-even combination is associated with the *SM* $SU(2)_L$ doublet, while the T-odd combination is its T-parity partner. The T-odd fermion sector consists of three generations of the mirror quarks and leptons with vectorial couplings under $SU(2)_L \times U(1)_Y$. Only T-odd leptons are related our calculation, we denote them by

$$\begin{pmatrix} \nu_H^1 \\ \ell_H^2 \end{pmatrix}, \quad \begin{pmatrix} \nu_H^2 \\ \ell_H^2 \end{pmatrix}, \quad \begin{pmatrix} \nu_H^3 \\ \ell_H^3 \end{pmatrix} \quad (1)$$

with their masses satisfying to first order in ν/f [10]

$$M_{\nu_H}^1 = M_{\ell_H}^1, \quad M_{\nu_H}^2 = M_{\ell_H}^2, \quad M_{\nu_H}^3 = M_{\ell_H}^3. \quad (2)$$

Here $\nu = 246 GeV$ is the electroweak scale and f is the scale parameter of the gauge symmetry breaking of the *LHT* model. There are $M_i^i = \sqrt{2}k_i^i f$, in which k_i^i are the eigenvalues of the mass matrix k and their values are generally dependent on the lepton species i . The coupling expressions of the T-odd lepton to other particles are given in Refs.[7,10].

It is well known that, at the *LHC*, the gluon distribution function is considerably larger than that for the quark. So we first consider the gluon-induced production of the T-odd lepton pairs. In order to produce the lepton pairs from gluon fusion mechanism, we must have a virtual quark triangular loop, which is connected to the lepton pairs by

a neutral boson (photon γ , gauge boson Z , or scalar H). However, the contributions of the photon vector coupling and the Z vector coupling vanish due to Furry's theorem [3]. From Refs.[7,10], one can see that the Z axial-vector coupling to the T-odd leptons is equal to zero and the scalar H can not couple to the T-odd quarks T_- and D_H . Thus, the T-odd lepton pair can only be produced via the s -channel H exchange diagram, in which the quark triangular loops include the top quark t , the up-type T-odd quark U_H , and the T-even vectorlike quark T_+ .

Using the relevant Feynman rules given in Refs.[7,10], we can easily calculate the hadronic production cross section σ_g of the gluon fusion process $gg \rightarrow H \rightarrow \ell_H^+ \ell_H^-$ at the LHC , which depend on the T-odd quark masses, the T-odd lepton masses, the mixing parameter x_L , and the scale parameter f . The mass of the T-even vectorlike top quark T_+ can be determined by the free parameters x_L and f . To simplify our calculation, we will take the free parameter $x_L = 0.5$ and the *Higgs* boson mass $M_H = 120 GeV$. For the T-odd quark masses, we will take $M_{U_H}^1 = M_{U_H}^2 = M_{U_H}^3 = M_U$ and assume that their values are smaller than $4 TeV$. M_ℓ is the T-odd lepton mass taken as $M_{\ell_H}^1 = M_{\ell_H}^2 = M_{\ell_H}^3 = M_\ell$ and assumed that its value is in the range of $200 GeV \sim 1000 GeV$. Recently, the improved parton distribution functions (*PDFs*) have been given in the literature, for example in Refs.[13,14]. However, as numerical estimation, we will use *CTEQ6L PDF*[15] for the gluon *PDF* and assume that the renormalization scale μ_R and the factorization scale μ_F have the relation $\mu_F = \mu_R = 2M_\ell$. Our calculation results show that the production cross section σ_g is very small and its value is smaller than $1 fb$ in most of the parameter space of the *LHT* model. This is because the coupling of the *Higgs* boson H to the T-odd leptons is proportion to the factor ν^2/f^2 and the production cross section σ_g is suppressed at least by the factor ν^4/f^4 .

The T-odd lepton pairs can also be produced via the *Drell – Yan* process and the $\gamma\gamma$ fusion mechanism. The former process is induced by the s -channel γ exchange and Z exchange, while the latter process can proceed via the t -channel and u -channel T-odd lepton exchanges. Since the $H\gamma\gamma$ and $H\ell_H^+ \ell_H^-$ couplings are very small, we will ignore the contributions of the s -channel process $\gamma\gamma \rightarrow H \rightarrow \ell_H^+ \ell_H^-$. In this case, it is obvious that

the production cross sections of the subprocesses $q\bar{q} \rightarrow \ell_H^+ \ell_H^-$ and $\gamma\gamma \rightarrow \ell_H^+ \ell_H^-$ are only dependent on the model dependent parameter — the T-odd lepton mass M_ℓ .




Figure 1: The production cross sections σ_q (a) and σ_γ (b) as function of the T-odd lepton mass M_ℓ .

Using the equivalent photon approximation (EPA) approach [16], the hadronic cross section σ_γ at the *LHC* can be obtained by folding the cross section for the subprocess $\gamma\gamma \rightarrow \ell_H^+ \ell_H^-$ with the photon distribution function $f_{\gamma/p}$, which can be written as $f_{\gamma/p} = f_{\gamma/p}^{el} + f_{\gamma/p}^{inel}$. $f_{\gamma/p}^{el}$ and $f_{\gamma/p}^{inel}$ are the elastic and inelastic components of the equivalent photon distribution of the proton, which have been extensively studied in Refs.[17, 18]. Our numerical results are given in Fig.1, in which the pair production cross sections σ_q (Fig.1a) and σ_γ (Fig.1.b) are plotted as functions of the T-odd mass M_ℓ . In our numerical calculation, we have assumed that the photon distribution function $f_{\gamma/p}$ includes both the elastic and inelastic components of the equivalent photon distribution of the proton and have taken the *CTEQ6L PDFs* for the quark distribution functions $f_{q_i/p}$. One can see from Fig.1 that the production cross section of the T-odd lepton pairs induced by the *Drell – Yan* process is significantly larger than that for the $\gamma\gamma$ fusion mechanism. For $200\text{GeV} \leq M_\ell \leq 1000\text{GeV}$, the values of the cross sections σ_q and σ_γ are in the ranges

of $270.6fb \sim 0.3fb$ and $26.4fb \sim 0.11fb$, respectively. If we assume the yearly integrated luminosity $\mathcal{L}_{int} = 100fb^{-1}$ for the *LHC* with the *c.m.* energy $\sqrt{s} = 14TeV$, then there will be several up to ten thousands of the $\ell_H^+\ell_H^-$ events to be generated per year.

It is well known that one of the main tasks of the *LHC* is to determine whether the breaking of the electroweak symmetry is due to the *SM Higgs* boson. The vector boson fusion (*VBF*) mechanism is the second most copious source for the *SM Higgs* boson production at the *LHC*, which can be used to directly study the exact dynamics of the electroweak symmetry breaking. The new charged lepton pairs can also be produced via the *VBF* mechanism at the *LHC* [4]. Pair production of the T-odd leptons predicted by the *LHT* model can be induced by the *s*-channel processes $W^\pm W^\mp \rightarrow \gamma, Z, H \rightarrow \ell_H^+\ell_H^-$ and $ZZ \rightarrow H \rightarrow \ell_H^+\ell_H^-$, and can also proceed via the *t*-channel T-odd neutrino ν_H exchange and T-odd lepton ℓ_H exchange for the processes $W^\pm W^\mp \rightarrow \ell_H^+\ell_H^-$ and $ZZ \rightarrow \ell_H^+\ell_H^-$, respectively.




Figure 2: The production cross section σ_W for the subprocess $W^+W^- \rightarrow \ell_H^+\ell_H^-$ as a function of the T-odd lepton mass M_ℓ .

For $M_\ell > 200GeV$, the *c.m.* energy $\sqrt{\hat{s}}$ of the subprocesses $W^\pm W^\mp \rightarrow \ell_H^+\ell_H^-$ is larger

than 400GeV and there is $\hat{s} \gg M_W^2$. In this energy region, the gauge bosons $W^\pm$ coming from quark or antiquark can be treated as on-shell. So, in our estimating the hadronic production cross section of the subprocesses $W^\pm W^\mp \rightarrow \ell_H^+ \ell_H^-$, we will adopt the effective W -boson approximation (EWA) method [19] and include all contributions of the longitudinal and transverse W -boson components. The structure functions for the gauge bosons $W^\pm$ are taken as the forms given by Ref.[20], in which authors have shown that the EWA method approximates very well the exact result. Considering the coupling of the *Higgs* boson H to the T-odd lepton pairs, which can be approximately written as $-\frac{iM_\ell}{\nu} \frac{\nu^2}{4f^2}$, is very small, we have neglected the contributions of H exchange. Our numerical results are shown in Fig.2. One can see from Fig.2 that the value of the production cross section can only reach 2.5fb in most of the parameter space, which is smaller than that for the $\gamma\gamma$ fusion mechanism.

Similar with that for the processes $W^\pm W^\mp \rightarrow \ell_H^+ \ell_H^-$, we can also neglect the contributions of the s -channel H exchange to the process $ZZ \rightarrow \ell_H^+ \ell_H^-$. Thus, this process is mainly induced by the t -channel and u -channel T-odd lepton exchanges. It is well known that the structure function for the gauge boson Z is smaller than that for the gauge boson W , the hadronic cross section for the subprocess $ZZ \rightarrow \ell_H^+ \ell_H^-$ is suppressed at least by an order of magnitude compared to that for the subprocess $W^+ W^- \rightarrow \ell_H^+ \ell_H^-$. For $200\text{GeV} \leq M_\ell \leq 1000\text{GeV}$, its value is in the range of $1.35 \times 10^{-1}\text{fb} \sim 2.5 \times 10^{-3}\text{fb}$. We further estimate the hadronic cross section for the subprocess $Z\gamma \rightarrow \ell_H^+ \ell_H^-$, our numerical results show that its value is in the range of $1.25\text{fb} \sim 6.9 \times 10^{-3}\text{fb}$ for $200\text{GeV} \leq M_\ell \leq 1000\text{GeV}$.

3. Conclusions and discussions

In this letter, we consider various possible production channels of the T-odd lepton pairs at the *LHC*. We find that the most important production channel for the T-odd lepton pairs predicted by the *LHT* model is the *Drell – Yan* process, i.e. $q\bar{q} \rightarrow \ell_H^+ \ell_H^-$. For $200\text{GeV} \leq M_\ell \leq 1000\text{GeV}$, the value of the hadronic production cross section is in the range of $270.6\text{fb} \sim 0.3\text{fb}$. The second important production channel is the $\gamma\gamma$ fusion mechanism, i.e. $\gamma\gamma \rightarrow \ell_H^+ \ell_H^-$. With reasonable values of the free parameters for the *LHT*

model, the production cross section value can reach $26.4fb$.

The T-odd leptons are always heavier than the T-odd gauge boson B_H . They are lighter or heavier than the T-odd gauge bosons W_H and Z_H , which depend on the values of the coupling parameter $k_l = k_l^1 = k_l^2 = k_l^3$. For $k_l < 0.45$, there is $M_l < M_{Z_H} \approx M_{W_H}$. Thus, for $k_l > 0.45$, the possible decay modes of the T-odd lepton ℓ_H are $B_H l$, $Z_H l$, and $W_H \nu$. However, as long as $k_l \leq 1$, ℓ_H mainly decays to $B_H \ell$ and there is $Br(\ell_H \rightarrow B_H \ell) \simeq 100\%$ for $k_l < 0.45$. If we assume that the T-odd lepton ℓ_H decays to $B_H \ell$, then pair production process of the T-odd leptons at the *LHC*, i.e. the process $PP \rightarrow \ell_H^+ \ell_H^- + X$, can give rise to the $\overline{B}_H B_H l^+ l^-$ final state. The new gauge boson B_H predicted by the *LHT* model is the stable and lightest T-odd particle, which can be seen as an attractive dark matter candidate [7]. Certainly, if the T-parity is violated by anomalies, the T-odd gauge boson B_H can decay into the *SM* gauge boson pairs WW and ZZ [21,22]. If we assume that T-parity is strictly conserved and the T-odd gauge boson B_H can be seen as missing energy, then pair production of the T-odd leptons at the *LHC* can induce the opposite-sign same-flavor leptons plus missing energy ($\ell^+ \ell^- + \cancel{E}$) signature. The main backgrounds for this kind of signals come from the *SM* processes $PP \rightarrow ZZ + X$, $PP \rightarrow \tau\tau + X$, and $PP \rightarrow W^+W^- + X$, in which the lepton τ and the gauge bosons Z and W decay leptonically. Detailed analysis of the signals and backgrounds have been given in Ref.[23]. They have shown that the ZZ background can be eliminated by calculating the invariant mass of the charged lepton pair, which can reconstruct the Z boson mass. It is possible to extract the $\ell^+ \ell^- + \cancel{E}$ signal from the $\tau\tau$ background by applying suitable cuts. The W^+W^- background is much large, so it is very challenging to extract the $\ell^+ \ell^- + \cancel{E}$ signal from this background. However, it has been shown that the appropriate cuts can significantly suppress the W^+W^- background [24]. Certainly, detailed confirmation of the observability of the $\ell^+ \ell^- + \cancel{E}$ signal, which comes from the subprocess $q\bar{q} \rightarrow \ell_H^+ \ell_H^-$, would require Monte-Carlo simulations of the signal and background.

Although the contributions of the $\gamma\gamma$ fusion mechanism to pair production of the T-odd lepton pairs at the *LHC* are much smaller than those for the *Drell – Yan* process.

For $M_\ell \geq 200\text{GeV}$, its production cross section can only reach 26.4fb . However, due to absence of the proton remnants, this production process has clean experimental conditions and well defined final states, which can be selected and precisely reconstructed [25]. The SM backgrounds coming from the partonic interactions, such as $PP \rightarrow W^+W^-$, ZZ , and $\tau\bar{\tau}$, can be significantly omitted by using the large rapidity gap technique and the dedicated very forward detectors [26]. The irreducible backgrounds, such as $\gamma\gamma \rightarrow W^+W^-$ and $\gamma\gamma \rightarrow \tau\bar{\tau}$ can be largely suppressed by applying acceptance cuts. Certainly, detailed detector simulation studies are needed.

Acknowledgments

This work was supported in part by the National Natural Science Foundation of China under Grants No. 10675057, Specialized Research Fund for the Doctoral Program of Higher Education (SRFDP) (No. 200801650002), the Natural Science Foundation of the Liaoning Scientific Committee (No. 20082148), and Foundation of Liaoning Educational Committee (No. 2007T086).

References

- [1] F. M. L. Almeida et al., *Phys. Rev. D***44**(1991)2836; G. Azuelos, A. Djouadi, Z. *Phys. C***63**(1994)327; F. M. L. Almeida et al., *Phys. Rev. D***51**(1995)5990; F. M. L. Almeida et al., *Phys. Rev. D***63**(2001)075005; F. M. L. Almeida et al., *Eur. Phys. J. C***30**(2003)327; A. T. Alan, A. T. Tascai, N. Kavagoz, *Mod. Phys. Lett. A***21**(2006)1869.
- [2] W. Buchmuller, C. Greub, *Nucl. Phys. B***363**(1991)345; Y. A. Coutinho, A. J. Rasmalho, S. Wulck, *Phys. Rev. D***54**(1996)1215; F. M. L. de Almeida et al., *Phys. Rev. D***65**(2002)115010; A. T. Alan, A. T. Tascai, O. Cakir, *Acta. Phys. Polon B***35**(2004)2199; A. T. Tascai, A. T. Alan, *hep-ph/0608045*.

- [3] S. S. D. Willenbrock, D. A. Dicus, *Phys. Lett. B***156**(1985)429; J. E. C. Montalvo, O. J. P. Eboli, S. F. Novaes, *Phys. Rev. D***46**(1992)181; P. H. Frampton, D. Ng, M. Sher, Y. Yuan, *Phys. Rev. D***48**(1993)3128; Y. D. A. Coutinho, J. A. M. Simoes, C. M. Porto, P. P. Q. Filho, *Phys. Rev. D***57**(1998)6975.
- [4] O. J. P. Eboli, G. C. Marques, S. F. Novces, A. A. Natale, *Phys. Rev. D***34**(1986)771; S. Darvson, S. S. D. Willenbrock, *Nucl. Phys. B***284**(1987)449; D. A. Dicus, *Nucl. Phys. B***287**(1987)397; C. P. Yuan, *Nucl. Phys. B***310**(1988)1.
- [5] M. Drees, R. M. Godbole, M. Nowakowski, S. D. Rindani, *Phys. Rev. D* **50**(1994)2335; G. Bhattacharya, P. Kalyniak, K. A. Peterson, *Phys. Rev. D***53** (1996)2371; M. M. Boyce, M. A. Doncheski, H. Konig, *Phys. Rev. D***55**(1997)68.
- [6] H. C. Cheng, I. Low, *JHEP* **0309**(2003)051; *JHEP* **0408**(2004)061; I. Low, *JHEP* **0410**(2004)067.
- [7] J. Hubisz, P. Meade, *Phys. Rev. D***71**(2005)035016; J. Hubisz, P. Meade, A. Noble and M. Perelstein, *JHEP* **0601**(2006)135; C. S. Chen, Kingman Cheung and T. -C. Yuan, *Phys. Lett. B***644**(2007)158; C. -R. Chen, K. Tobe, C. -P. Yuan, *Phys. Lett. B***640**(2006)263.
- [8] A. Belyaev, Chuan-Ren Chen, K. Tobe, C. P. Yuan, *Phys. Rev. D* **74**(2006)115020; A. Freitas, D. Wyler, *JHEP* **0611**(2006)061.
- [9] Q. H. Cao, C. S. Li, C. P. Yuan, *Phys. Lett. B***668**(2008)24; M. Blanke et al, *JHEP* **0612**(2006)003; D. Choudhury, D. K. Ghosh, *JHEP* **0708**(2007)084; S. Matsumoto, M. M. Nojiri, D. Nomura, *Phys. Rev. D***75** (2007)055006; K. Kong, S. C. Park, *JHEP* **0708**(2007)038.
- [10] M. Blanke et al., *JHEP* **0701**(2007)066; F. del Aguila, J. I. Illana, M.D. Jenkins, *JHEP* **0901**(2009)080.

- [11] Lei Wang, Wen-Yu Wang, Jin-Min Yang, Huan-Jun Zhang, *Phys. Rev. D* **76**(2007)017702; Xue-Lei Wang, Yan-Yu Zhang, Hui-Ling Jin, Yan-Hui Xi, *Nucl. Phys. B* **810**(2009)226.
- [12] Chong-Xing Yue, Nan Zhang, Shi-Hai Zhu, *Eur. Phys. J. C* **53**(2008)215.
- [13] W. K. Tung et al., *JHEP* **0702**(2007)053; H. L. Lai et al., *JHEP* **0704**(2007)089; J. Pumplin, H. L. Lai and W. K. Tung, *Phys. Rev. D* **75**(2007)054029.
- [14] P. M. Nadolsky et al., *Phys. Rev. D* **78**(2008)013004.
- [15] J. Pumplin et al. (CTEQ Collaboration), *JHEP* **0602** (2006)032.
- [16] V. M. Budnev, I. F. Ginzburg, G. V. Meledin, and V. G. Serbo, *Phys. Rept.* **15**(1974)181.
- [17] M. Glück, C. Pisano and E. Reya, *Phys. Lett. B* **540**(2002)75; C. Pisano, *Eur. Phys. J. C* **38**(2004)79, and reference therein.
- [18] G. Altarelli, G. Martinelli, B. Mele and R. Rückl, *Nucl. Phys.* **B262**(1985)204; G. Martinelli, B. A. Kniehl, *Phys. Lett. B* **254**(1991)267; A. Mukherjee and C. Pisano, *Eur. Phys. J. C* **30**(2003)477.
- [19] M. S. Chanowitz and M. K. Gaillard, *Phys. Lett. B* **142** (1984)85; G. L. Kane, W. W. Repko and W. B. Rolnick, *Phys. Lett. B* **148**(1984)367; S. Dawson, *Nucl. Phys. B* **249**(1985)42; J. Lindfors, *Z. Phys. C* **28**(1985)427; J. F. Gunion, J. Kalinowski and A. Tofghi-Niaki, *Phys. Lett.* **57**(1986)2351.
- [20] A. Alboteanu, W. Kilian, and J. Reuter, *JHEP* **0811**(2008)010.
- [21] C. T. Hill and R. J. Hill, *Phys. Rev. D* **75**(2007)115009; *D* **76**(2007)115014.
- [22] V. Barger, Wai-Yee Keung, Y. Gao, *Phys. Lett. B* **655** (2007)228; D. Krohn, I. Yavin, *JHEP* **0806**(2008)092; A. Freitas, P. Schwaller, D. Wyler, *JHEP* **0809**(2008)013.

- [23] V. D. Barger, Tao Han, J. Ohnemus, *Phys. Rev. D***37** (1988)1174; I. Hinchliffe, *Int. J. Mod. Phys. A***4**(1989)3867.
- [24] M. Dittmar and H. K. Dreiner, *Phys. Rev. D***55**(1997)167; G. Davatz, M. Dittmar and A. S. Giolo-Nicollerat, *J. Phys. G***33**(2007)N85.
- [25] M. Klasen, *Nucl. Phys. B, Proc. Suppl.* **179-180**(2008)298; M. V. Donckt, *arXiv:0806.4865[hep-ph]*; K. Piotrkowski, *Nucl. Phys. B, Proc. Suppl.* **179-180**(2008)2.
- [26] N. Schul and K. Piotrkowski, *Nucl. Phys. B, Proc. Suppl.* **179-180**(2008)289; M. Boonekamp, J. Cammin, R. Peschanski, C. Royon, *Phys. Lett. B***654**(2007)104; T. Pilerzchala and K. Piotrkowski, *Nucl. Phys. B, Proc. Suppl.* **179-180**(2008)257; O. Kepka and C. Royon, *Phys. Rev. D***78**(2008)073005; S. Oryn, *Nucl. Phys. B, Proc. Suppl.* **179-180**(2008)269; J. de Favereau de Jeneret, S. Oryn, *Nucl. Phys. B, Proc. Suppl.* **179-180**(2008)277.